

Graded Lie algebras with finite polydepth

Y. Felix, S. Halperin and J.-C. Thomas

February 1, 2008

Abstract

If A is a graded connected algebra then we define a new invariant, polydepth A , which is finite if $\text{Ext}_A^*(M, A) \neq 0$ for some A -module M of at most polynomial growth. **Theorem 1:** If $f : X \rightarrow Y$ is a continuous map of finite category, and if the orbits of $H_*(\Omega Y)$ acting in the homology of the homotopy fibre grow at most polynomially, then $H_*(\Omega Y)$ has finite polydepth. **Theorem 5:** If L is a graded Lie algebra and polydepth UL is finite then either L is solvable and UL grows at most polynomially or else for some integer d and all r , $\sum_{i=k+1}^{k+d} \dim L_i \geq k^r$, $k \geq$ some $k(r)$.

AMS Classification : 55P35, 55P62, 17B55

Key words : Graded Lie algebras, loop space homology, depth, solvable Lie algebras.

We work over a field $\mathbb{k}$ of characteristic different from 2. If $V = \{V_k\}$ is a graded vector space we denote by $V^\# = \{\text{Hom}_{\mathbb{k}}(V_k, \mathbb{k})\}$ the dual graded vector space. A graded Lie algebra is a graded vector space L , equipped with a bilinear map $[\cdot, \cdot] : L_i \times L_j \rightarrow L_{i+j}$ satisfying

$$[x, y] + (-1)^{ij}[y, x] = 0$$

and

$$[x, [y, z]] = [[x, y], z] + (-1)^{ij}[y, [x, z]]$$

for $x \in L_i, y \in L_j, z \in L$. It follows that $3[x, [x, x]] = 0$ for x of odd degree, and so if $\text{char } k = 3$ we further require that $[x, [x, x]] = 0$. Finally we consider only graded Lie algebras satisfying $L = \{L_i\}_{i \geq 1}$ and each L_i is finite dimensional. (Any graded vector space V with each V_i finite dimensional is said to have *finite type*.)

The universal enveloping algebra of L is denoted by UL and it satisfies the classical Poincaré Birkhoff Witt Theorem (in characteristic 3 this uses the $[x, [x, x]] = 0$ requirement).

Important examples appear in topology. Let X be a simply connected topological space with rational homology of finite type. Then the rational homotopy Lie algebra L_X of X is defined by

$$L_X = \pi_*(\Omega X) \otimes \mathbb{Q}; \quad [\cdot, \cdot] = \text{Samelson product},$$

and the Hurewicz map extends to an isomorphism, [10],

$$UL_X \xrightarrow{\cong} H_*(\Omega X; \mathbb{Q}).$$

Analogously, if X is a finite n -dimensional and r -connected CW complex, then for primes $p > n/r$, $H_*(\Omega X; F_p) = UE$ for some graded Lie algebra E [8].

If M is a module over a (graded) algebra A then the *grade* of M , $\text{grade } M$, is the least integer q (or ∞) such that $\text{Ext}_A^q(M, A) \neq 0$. And if $V = \{V_i\}_{i \geq 0}$ is a graded vector space then V has *at most polynomial growth* if for some constant C , and some non-negative integer, d , $\sum_{i \leq n} \dim V_i \leq Cn^d$, $n \geq 1$. In this case the least such d is called the *polynomial bound* for the growth of V and is denoted by $\text{polybd}(V)$. If V does not have at most polynomial growth we put $\text{polybd } V = \infty$ and we say that V grows faster than any polynomial.

In this paper we combine these two notions in the

Definition: the *polygrade* of an A -module, M , is the sum, $\text{grade } M + \text{polybd } M$, and the *polydepth* of A is the least integer (or ∞) occurring as the polygrade of an A -module.

In the case $A = UL$ the unique augmentation $UL \rightarrow \mathbb{k}$ makes $\mathbb{k}$ into a UL -module, and by definition, the grade of $\mathbb{k}$ is the *depth* of UL . Since $\text{polybd } \mathbb{k} = 0$ it follows that:

$$\text{polydepth } UL \leq \text{depth } UL \quad (1)$$

Moreover (cf. Proposition 2) if $\dim L < \infty$ then equality holds. We shall abuse notation and refer to these invariants respectively as $\text{polydepth } L$ and $\text{depth } L$.

Note that $\text{Ext}_{UL}^0(UL, UL)$ contains the identity map and so

$$\text{polydepth } L \leq \text{polybd } UL. \quad (2)$$

Observe as well that for any graded vector space M , $\text{polybd } M = 0$ if and only if $\dim M$ is finite. Thus $\text{polydepth } L = 0$ if and only if $\text{depth } L = 0$, which happens if and only if L is finite dimensional and concentrated in odd degrees.

Depth has been a useful concept in topology because, on the one hand, Lusternik-Schnirelmann category satisfies [1]

$$\text{depth } L_X \leq \text{cat } X$$

and, on the other hand ([3], [4], [5]), finite depth has important implications for the structure of a graded Lie algebra.

The purpose of this paper is to show that essentially the same implications follow from the weaker hypothesis that $\text{polydepth } L$ is finite, while simultaneously identifying a larger class of topological spaces and Lie algebras for which the weaker hypothesis holds.

Indeed, we have

Theorem 1. *If $F \rightarrow X \xrightarrow{f} Y$ is a fibration of path-connected spaces, then*

$$\text{polydepth } H_*(\Omega Y) \leq \text{polybd } H_*(F) + \text{cat } f.$$

proof: The fibration determines an action up to homotopy of ΩY on F , which makes $H_*(F)$ into an $H_*(\Omega Y)$ -module. According to [6], $\text{grade } H_*(F) \leq \text{cat } f$. $\square$

Our main structural theorems read:

Theorem 2. *Let $E(L)$ denote the linear span of elements $x \in L_{\text{even}}$ such that adx acts nilpotently on each $y \in L$. Then*

$$\dim E(L) \leq \text{polydepth } L.$$

Theorem 3. *The following conditions on a graded Lie algebra L are equivalent*

- (i) L is the union of solvable ideals and polydepth L is finite;
- (ii) UL grows at most polynomially (polybd UL is finite);
- (iii) L_{even} is finite dimensional, and for some constant C

$$\sum_{i \leq n} \dim L_i \leq C \log_2 n, \quad n \geq 1.$$

In this case L is solvable.

Theorem 4. *If L is a graded Lie algebra of finite polydepth then the union of the solvable ideals of L is a solvable ideal of finite polydepth.*

Theorem 5. *Suppose polydepth L is finite and L is not solvable. Then there is an integer d such that for all $r \geq 1$:*

$$\sum_{i=k+1}^{k+d} \dim L_i \geq k^r, \quad k \geq \text{some } k(r).$$

Remark. In [7] it is shown that if $L = L_X$ where X is a finite 1-connected CW complex, then we may take $d = \dim X$ in Theorem 5.

1 Properties of polydepth

Lemma 1: *If M is a module for some graded algebra A of finite type and if $\text{Ext}_A^q(M, A) \neq 0$ then $\text{Ext}_A^q(A \cdot x, A) \neq 0$ for some x in a subquotient module of M .*

proof: Recall that $A^\# = \text{Hom}_{\mathbb{k}}(A, \mathbb{k})$. Then $\text{Ext}_A^q(M, A)$ is the dual of $\text{Tor}_q^A(M, A^\#)$ and a direct limit argument shows that for some $x_1, \dots, x_n \in M$,

$$\text{Tor}_q^A(A \cdot x_1 + \dots + A \cdot x_n, A^\#) \neq 0.$$

Now use the exact sequence associated to the inclusion $A \cdot x_1 + \dots + A \cdot x_{n-1}$ in $A \cdot x_1 + \dots + A \cdot x_n$. $\square$

Corollary: Polydepth A is the least m such that polygrade $N = m$ for some monogenic A -module N .

Remark: It follows from the Corollary that we may improve Theorem 1 to the inequality

$$\text{polydepth } H_*(\Omega Y) \leq \text{polybd}(H_*(\Omega Y) \cdot \alpha) + \text{cat } f, \quad \text{some } \alpha \in H_*(F). \quad (3)$$

Proposition 1. *Let L be a graded Lie algebra.*

- (i) *Each ideal satisfies polydepth $I \leq \text{polydepth } L$.*
- (ii) *Let E be a Lie subalgebra of L . If L has finite polydepth and if for each $x \in L/E$ the orbit $UE \cdot x$ has at most polynomial growth, then E has finite polydepth.*
- (iii) *For n sufficiently large the sub Lie algebra E generated by $L_{\leq n}$ satisfies polydepth $E \leq \text{polydepth } L$.*

proof: (i) This follows from the Hochschild-Serre spectral sequence, converging from $\text{Ext}_{UL/I}^p(\mathbb{k}, \text{Ext}_{UI}^q(M, UL))$ to $\text{Ext}_{UL}^{p+q}(M, UL)$. (Note that since UL is UI -free, $\text{grade}_{UI}(M)$ is the least q such that $\text{Ext}_{UI}^q(M, UL) \neq 0$.)

(ii) As in Lemma 1, $\text{Ext}_{UL}^q(M, UL)$ is dual to $\text{Tor}_q^{UL}(M, (UL)^\#)$, and this is the homology of the Cartan-Eilenberg-Chevalley complex $\wedge^p sL \otimes M \otimes (UL)^\#$. Write $L = E \oplus V$ and set $F_p = \wedge^p sE \otimes \wedge^{\leq p} sV \otimes M \otimes (UL)^\#$. This filtration determines a convergent spectral sequence, introduced by Koszul in [9], and which is the Hochschild-Serre spectral sequence when E is an ideal. The E^1 -term of the spectral sequence is $\text{Tor}_q^{UE}(\wedge^p sL/E \otimes M, (UL)^\#)$, converging to $\text{Tor}_{p+q}^{UL}(M, (UL)^\#)$.

Each element $z \in \wedge^p sL/E \otimes M$ is contained in a finite sum of UE -modules of the form $s(UE \cdot x_1) \wedge \cdots \wedge s(UE \cdot x_p) \otimes M$ and it follows that

$$\text{polybd}(UE \cdot z) \leq p \cdot \text{polybd}(UE \cdot x) + \text{polybd}(M)$$

for some $x \in L/E$. Choose M so that $\text{polydepth } L = \text{polygrade } M$ and apply Lemma 1 with $p + q = \text{grade } M$.

(iii) If $\text{Ext}_{UL}^p(M, UL)$ is non-zero and $\text{polybd}(M) < \infty$ it suffices to choose E so that the restriction $\text{Ext}_{UL}^p(M, UL) \rightarrow \text{Ext}_{UE}^p(M, UL)$ is non zero. $\square$

Corollary 1. (of the proof of (ii)). Suppose for some $k \geq 1$ that $\text{polybd}(UE \cdot x) \leq k$, $x \in L/E$. Then $\text{polydepth } E \leq k \text{ polydepth } L$.

Corollary 2. Let E be a sub-Lie algebra of a graded Lie algebra L . If L has finite polydepth and L/E has at most polynomial growth, then E has finite polydepth.

Example 1. Let $\mathbb{L}(V)$ be the free Lie algebra on a graded vector space V . Then for any graded Lie algebra L , $L \coprod \mathbb{L}(V)$ has depth 1. Thus the injection $L \rightarrow L \coprod \mathbb{L}(V)$ shows that each graded Lie algebra is a sub-Lie algebra of a Lie algebra of finite polydepth. The previous corollary gives restriction on a Lie algebra L for being a sub-Lie algebra of a Lie algebra of finite polydepth, K , when the quotient has at most polynomial growth.

Proposition 2. If L is a finite dimensional graded Lie algebra then

$$\text{polydepth } L = \text{depth } L.$$

proof: As observed in the introduction, $\text{polydepth } L \leq \text{depth } L$. On the other hand, by Lemma 1, $\text{polydepth } L = \text{polygrade } M$ for some monogenic module $M = UL \cdot x$. Now Theorem 3.1 in [3] asserts that $\text{polygrade } M = \text{depth } L$. $\square$

2 Proof of Theorem 2

Suppose $I \subset L$ is an ideal. If $\text{Ext}_{UL}^m(M, UL) \neq 0$, then $\text{Ext}_{UL/I}^p(\text{Tor}_q^{UI}(M, \mathbb{k}), UL/I) \neq 0$, some $p + q = m$. (Same proof as in: [3], Lemma 4.3, for the case $M = \mathbb{k}$). By Lemma 1 there is a monogenic UL/I -module N such that N is a subquotient of $\text{Tor}_q^{UI}(M, \mathbb{k})$, and $\text{grade } N \leq p$.

Now suppose L/I is finite dimensional. Then Theorem 3.1 in [3] asserts that

$$\text{grade } N + \text{polybd } N = \dim (L/I)_{\text{even}}.$$

On the other hand, write $(L/I)_{\text{even}} = V \oplus W$ where V is the image of $E(L)$. Let $x_i \in L_{\text{even}}$, $y_j \in L_{\text{odd}}$ and $z_k \in E(L)$ represent respectively bases of W , $(L/I)_{\text{odd}}$ and V . Then the elements $x_1^{k_1} \cdots x_s^{k_s} y_1^{\varepsilon_1} \cdots y_t^{\varepsilon_t} z_1^{m_1} \cdots z_u^{m_u}$, where $\varepsilon_i = 0$ or 1, represent a basis for UL/I .

Choose the z_k to act locally nilpotently in L . Then this basis applied to any $\omega \in \wedge^q sI$, shows that $\text{polybd}(UL/I) \cdot \omega \leq \dim W$. Hence if $u \in \wedge^q sL \otimes M$ represents a generator of N then

$$\text{polybd } N \leq \text{polybd}(UL/I \cdot u) \leq \text{polybd } M + \dim W.$$

Substitution in the equation above gives

$$\begin{aligned} \dim(L/I)_{\text{even}} &\leq \text{grade } N + \text{polybd } M + \dim W \\ &\leq \text{grade } M + \text{polybd } M + \dim W. \end{aligned}$$

Choose M so $\text{grade } M + \text{polybd}(M) = \text{polydepth } L$ and choose $I = L_{>2k}$. Then $V \cong E(L)_{\leq 2k}$ and we have

$$\dim E(L)_{\leq 2k} \leq \text{polydepth } L.$$

Since this holds for all k the theorem is proved. $\square$

3 Solvable Lie algebras

Lemma 2. *Let L be a Lie algebra concentrated in odd degrees. Then $\text{Ext}_{UL}(-, UL) = \text{Hom}_{UL}(-, UL)$. In particular*

$$\text{polydepth } L = \text{polybd } UL.$$

proof: Since $L = L_{\text{odd}}$ it is necessarily abelian. Now $\text{Ext}_{UL}(-, UL)$ is the dual of $\text{Tor}^{UL}(-, (UL)^{\#})$ and this is the limit of $\text{Tor}^{UL_{\leq n}}(-, (UL)^{\#})$, which dualizes to $\text{Ext}_{UL_{\leq n}}(-, UL)$. Since $UL_{\leq n}$ is a finite dimensional exterior algebra and UL is $UL_{\leq n}$ -free it follows that $\text{Ext}_{UL_{\leq n}}^+(-, UL) = 0$, and so $\text{Ext}_{UL}^+(-, UL) = 0$.

Finally, since $\text{Ext}_{UL}^0(UL, UL)$ is non-zero, $\text{polydepth } L \leq \text{polybd } UL$. On the other hand if $\text{polydepth } L = m < \infty$, then for some M , we have $\text{Ext}_{UL}^p(M, UL) \neq 0$ and $\text{polybd } M = m - p$. By the above, $p = 0$ and so there is a non zero UL -linear map $f : M \rightarrow UL$. Any $f(m)$ is in some $UL_{<n}$ and if $f(m) \neq 0$ it follows that $UL_{\geq n} \xrightarrow{\cong} UL_{\geq n} \cdot m$. This implies $\text{polybd } M \geq \text{polybd } UL$ and $\text{polydepth } L \geq \text{polybd } UL$. $\square$

Lemma 3. *Let L be a graded Lie algebra of finite polydepth. If I is an ideal in L and $\text{polybd } I < \infty$ then $\text{polydepth } L/I < \infty$.*

proof: Choose M so that $\text{polygrade } M = \text{polydepth } L$. If $m = \text{grade } M$ then it follows (as in [3], proof of Theorem 4.1 for the case $M = k$) that for some p ,

$$\text{Ext}_{UL/I}^p(\text{Tor}_{m-p}^{UI}(M, k), UL/I) \neq 0.$$

Since $\text{Tor}_{m-p}^{UI}(M, k)$ is a subquotient of $\wedge^{m-p} sI \otimes M$ it follows that it has polynomial growth at most equal to $(m - p)\text{polybd } I$ $\square$

proof of Theorem 3:

(i) $\Rightarrow$ (ii). Let I be the sum of the ideals in L concentrated in odd degrees. Then I is an ideal of this form, necessarily abelian, and L/I has no ideals concentrated in odd degrees. Moreover $\text{polybd } UI = \text{polydepth } I \leq \text{polydepth } L$ (Lemma 2 and Proposition 1) and hence $\text{polydepth } L/I < \infty$ (Lemma 3).

Next we show that every solvable ideal J in L/I is finite dimensional, by induction on the solvlength. Indeed, if J is abelian then $J_{\text{even}} = E(J)$. Since $\text{polydepth } J \leq$

polydepth L/I (Proposition 1), Theorem 2 asserts that J_{even} is finite dimensional. Thus for some r , $J_{\geq r}$ is an ideal concentrated in odd degrees; i.e. $J_{\geq r} = 0$.

Now if J has solvlength k then its $(k+1)$ st derived algebra is abelian and so finite dimensional. Thus for some r , $J_{\geq r}$ has solvlength $k-1$. By induction, J is finite dimensional.

By hypothesis L/I is the sum of its solvable ideals. Since these are finite dimensional, each $x \in (L/I)_{\text{even}}$ acts locally nilpotently. Thus $(L/I)_{\text{even}} = E(L/I)$, and this is finite dimensional by Theorem 2. But $L_{\text{even}} \cong (L/I)_{\text{even}}$ since I is concentrated in odd degrees.

Suppose $L_{\text{even}} \subset L_{\leq 2n}$. Since $L_{>2n}$ is an ideal in odd degrees of finite polydepth, $\text{polybd } UL_{>n} < \infty$, while trivially $\text{polybd } UL/L_{>2n} < \infty$. Hence $\text{polybd } UL < \infty$.

(ii) $\Rightarrow$ (iii). Clearly $\text{polybd } UL \geq \dim L_{\text{even}}$, so the latter must be finite. It is trivial from the Poincaré Birkhoff Witt theorem that if $\sum_{i \leq n} \dim L_i = d(n)$ then

$$\sum_{i \leq nd(n)} \dim (UL)_i \geq 2^{d(n)}.$$

Thus $2^{d(n)} \leq K[nd(n)]^r$ for some constant K and some integer r , $r \geq 1$. It follows that $d(n) \leq \log_2 K + r \log_2 n + r \log_2 d(n) \leq r \log_2 n + \frac{1}{2}d(n)$, n sufficiently large.

(iii) $\Rightarrow$ (i). Choose N so that $I = L_{\geq N}$ is concentrated in odd degrees. Then UI is an exterior algebra and so

$$\sum_{i \leq n} \dim (UL)_i \leq 2^{\sum_{i \leq n} \dim L_i} \leq n^C$$

for some constant C . Thus, since L/I is finite dimensional, $\text{polybd } UL$ is finite. The identity of UL is in $\text{Ext}_{UL}^0(UL, UL)$ and so $\text{polydepth } L \leq \text{polybd } UL$.

Finally, since I is abelian and L/I is finite dimensional, L itself is solvable. This also proves the last assertion. $\square$

proof of Theorem 4: This is immediate from Proposition 1(i) and Theorem 3. $\square$

Proposition 3. Suppose I is a solvable ideal in a Lie algebra L of finite polydepth. Then

- (i) $\text{polydepth } L/I \leq \text{polydepth } L$
- (ii) $\dim I_{\text{even}} \leq \text{polydepth } I \leq \text{polybd } UI$.

proof: (i) As noted in the proof of Lemma 3,

$$\text{Ext}_{UL/I}^p(\text{Tor}_{m-p}^{UI}(M, k), UL/I) \neq 0,$$

where $m + \text{polybd } M = \text{polydepth } L$. Also the Tor is a subquotient of $\wedge^{m-p} sI \otimes M$. By Theorem 3

$$\sum_{i \leq n} \dim (\wedge^{m-p} sI \otimes M)_i \leq (C_1 \log_2 n)^{m-p} \cdot C_2 \cdot n^{\text{polybd } M}.$$

Hence $\text{polybd } \text{Tor}_{m-p}^{UI}(M, k) \leq \text{polybd } M + 1$, and so $\text{polydepth } L/I \leq \text{polydepth } M + p + 1$.

If $p < m$ then this gives $\text{polydepth } L/I \leq \text{polydepth } L$. If $p = m$ then $\text{Tor}_{m-p}^{UI}(M, k) = M \otimes_{UI} k$. Hence in this case $\text{polybd } \text{Tor}_{m-p}^{UI}(M, k) \leq \text{polybd } M$ and again $\text{polydepth } L/I \leq \text{polydepth } L$.

(ii) Since $I_{\text{even}} \subset I_{\leq 2n}$, some n (Theorem 3) we may apply the first assertion to obtain

$$\begin{aligned} \dim I_{\text{even}} &= \text{depth } I/I_{>2n} \text{ (cf. [2])} \\ &= \text{polydepth } I/I_{>2n} \text{ (Proposition 2)} \\ &\leq \text{polydepth } I. \end{aligned}$$

The second inequality has already been observed: $\text{polybd } UL = \text{polygrade } UL \geq \text{polydepth } L$. $\square$

Example 3. Consider a Lie algebra L concentrated in odd degrees with a basis $\{x_i \mid i \geq 1\}$ satisfying the degree relations

$$\deg x_i > \sum_{j < i} \deg x_j.$$

Then for each n , $\dim(UL)_n \leq 1$. The identity on UL shows that $\text{polydepth } L = 1$.

Example 4. Consider the graded Lie algebra $L = \mathbb{L}(a, x_n)_{n \geq 2}/I$, with $\deg a = 2$, $\deg x_n = 2^n + 1$, and where I is generated by the relations

$$[(\text{ada})^k x_r, (\text{ada})^l x_s] = 0, \quad k, l \geq 0, r, s \geq 2, \quad \text{and} \quad \text{ad}^{n+1}(a)(x_n) = 0.$$

Then $\text{polybd } UL = 2$, so that L has finite polygrade. On the other hand, L is solvable but not nilpotent, and is the union of the infinite sequence of the finite dimensional Lie algebras I_N generated by $a, x_2, \dots, x_N$.

Proposition 4. Let L be the direct sum of non-solvable Lie algebras $L(i)$, $i \leq n$. If $\text{polydepth } L(i) < \infty$ for $1 \leq i \leq n$, then

$$n \leq \text{polydepth } L \leq \sum_i \text{polydepth } L(i).$$

proof. We first prove by induction on n that for any UL -module M that has at most polynomial growth, we have $\text{Ext}_{UL}^{\leq n}(M, UL) = 0$. Consider the Hochschild-Serre spectral sequence

$$\text{Ext}_{UL(1)}^p(\text{Tor}_q^{U(L(2) \oplus \dots \oplus L(n))}(M, (U(L(2) \oplus \dots \oplus L(n))^{\#}), UL(1)) \Rightarrow \text{Ext}_{UL}^{p+q}(M, UL). \quad (4)$$

Since $L(1)$ commutes with the other $L(i)$ it follows that for each monogenic $UL(1)$ -module N that is a subquotient of $\text{Tor}_q^{U(L(2) \oplus \dots \oplus L(n))}(M, U(L(2) \oplus \dots \oplus L(n))^{\#})$ we have $\text{polybd } N \leq \text{polybd } M$.

Now since $L(1)$ is not solvable, $\text{polybd } UL(1) = \infty$ and the argument in the proof of Lemma 2 shows that $\text{Ext}_{UL(1)}^0(N, UL(1)) = 0$. Thus (Lemma 1) the left hand in (4) vanishes for $p = 0$. By induction on n it vanishes for $q < n - 1$ and so $\text{Ext}_{UL}^{\leq n}(M, UL) = 0$. Thus $\text{polydepth } L \geq n$.

On the other hand, there are $UL(i)$ -modules $M(i)$ such that $\text{polygrade } M(i) = \text{polydepth } L(i)$. Then $\otimes_{i=1}^n M(i)$ is a UL -module that has at most polynomial growth and whose polygrade is the sum of the polygrades of the $M(i)$. $\square$

4 Growth of Lie algebras

Proposition 5. Let L be a non solvable graded Lie algebra of finite polydepth. Then for each integer $r \geq 1$ there is a positive integer $d(r)$ such that

$$\sum_{i=k+1}^{k+d(r)} \dim L_i \geq k^r, \quad k \text{ sufficiently large.}$$

proof: We distinguish two cases.

Case A: L_{even} contains an infinite dimensional abelian sub Lie algebra E .

Choose n so that $\dim E_{\leq n} \geq (r+3)\text{polydepth } L$. Then there is a finite sequence

$$L = I(0) \supset I(1) \supset \cdots \supset I(l)$$

in which $I(j)$ is an ideal in $I(j-1)$ and $I(l)_{\leq n} = E_{\leq n}$.

By Proposition 1, $\text{polydepth } I(q) \leq \text{polydepth } L$. Thus without loss of generality we may suppose that $L = I(l)$; i.e. that $L_{\leq n}$ is an abelian sub Lie algebra concentrated in even degrees and that $\dim L_{\leq n} \geq (r+3)\text{polydepth } L$.

Let M be a UL -module such that $\text{grade } M + \text{polybd } M = \text{polydepth } L$ and put $m = \text{grade } M$. As observed in the proof of Proposition 1(ii), $\text{Ext}_{UL_{\leq n}}^q(\wedge^p sL_{>n} \otimes M, UL_{\leq n}) \neq 0$, for some $p+q = m$. It follows that for some $z \in \wedge^p sL_{>n} \otimes M$,

$$\text{polybd } UL_{\leq n} \cdot z + q \geq \dim L_{\leq n}$$

(Theorem 3.1 in [3]). Hence for some $x \in L$,

$$p(\text{polybd } UL_{\leq n} \cdot x) \geq \dim L_{\leq n} - q - \text{polybd } M.$$

Since $p+q+\text{polybd } M = \text{polydepth } L$ we conclude that

$$(2 + \text{polybd } UL_{\leq n} \cdot x) \cdot \text{polydepth } L \geq \dim L_{\leq n} \geq (r+3)\text{polydepth } L.$$

As observed in the introduction, since $\dim L = \infty$, $\text{polydepth } L > 0$. It follows that

$$\text{polybd } (UL_{\leq n} \cdot x) \geq r+1. \quad (5)$$

On the other hand, $UL_{\leq n}$ is the polynomial algebra $\mathbb{k}[y_1, \dots, y_s]$ on a basis $y_1, \dots, y_s$ of $L_{\leq n}$. Because of (5) it is easy to see (induction on s) that this basis can be chosen so that for some $w \in UL_{\leq n} \cdot x$, $\mathbb{k}[y_1, \dots, y_{r+1}] \rightarrow \mathbb{k}[y_1, \dots, y_{r+1}] \cdot w$ is injective. Put $d = \prod_{i=1}^{r+1} \deg y_i$ and note that

$$\sum_{i=k+\deg w+1}^{k+\deg w+d} \dim L_i \geq \sum_{i=k+1}^{k+d} \dim \mathbb{k}[y_1, \dots, y_{r+1}]_i \geq \frac{1}{r!} k^r.$$

It follows that for any r , and k sufficiently large,

$$\sum_{i=k+1}^{k+d} \dim L_i \geq k^{r-1}.$$

This proves Proposition 5 in case A.

Case B Every abelian sub Lie algebra of L_{even} is finite dimensional.

Let I be the sum of the solvable ideals in L . Then I_{even} is finite dimensional and $\text{polydepth } L/I$ is finite (Theorem 3 and Proposition 3). Thus all abelian sub Lie algebras of L/I are finite dimensional. Thus it is sufficient to prove Case B when L has no solvable ideals.

There are now two possibilities: either $L = L_{\text{even}}$, or L has elements of odd degree. In the latter case the sub Lie algebra generated by L_{odd} is an ideal, hence non-solvable and of finite polydepth. Let $L(s)$ denote the sub Lie algebra generated by the first s linearly independent elements $x_1, \dots, x_s$ of odd degree. For s sufficiently large, $\text{polydepth } L(s) \leq \text{polydepth } L$ (Proposition 1) and $\dim L(s)_{\text{even}} > \text{polydepth } L$ (obvious). Thus $L(s)$ cannot be solvable (Theorem 3). In other words, we may assume that either $L = L_{\text{even}}$ or else L is

generated by finitely many elements $x_1, \dots, x_s$ of odd degree. In either case set $E = L_{\text{even}}$, and note that $\dim E$ is infinite.

Define a sequence of elements z_i and sub Lie algebras $E(i)$ by setting $E(1) = E$, z_i is a non-zero element in $E(i)$ and $E(i+1) \subset E(i)$ is the sub Lie algebra of elements on which $\text{ad } z_i$ acts nilpotently.

Since E contains no infinite dimensional abelian Lie algebra some $E(N+1) = 0$ and $E(1)/E(2) \oplus \dots \oplus E(N)/E(N+1)$ is a graded vector space isomorphic with E .

Put $d = \prod \deg z_i$ and $d_i = d/\deg z_i$. Then $(\text{ad } z_1)^{d_1} \oplus \dots \oplus (\text{ad } z_N)^{d_N}$ is an injective transformation of $E(1)/E(2) \oplus \dots \oplus E(N)/E(N+1)$ of degree d . Since this space is isomorphic with E it follows that

$$\sum_{i=k+1}^{k+d} \dim E_i \geq \frac{d}{k+d} \sum_{i=1}^{k+d} \dim E_i, \quad k \geq 1. \quad (6)$$

On the other hand, choose n so that

$$\dim E_{\leq n} \geq (r+3) \cdot \text{polydepth } L.$$

(This is possible because E is infinite dimensional.) Set $I = L_{>n}$. Let M be an L -module with $\text{polydepth } L = \text{polygrade } M$. As in the proof of Theorem 4.1 in [3], $\text{Ext}_{UL/I}^p(\text{Tor}_q^{UI}(M, \mathbb{k}), UL/I) \neq 0$ for $p+q = \text{grade } M$. Thus p, q and $\text{polybd } M$ are all bounded above by $\text{polydepth } L$. Now Theorem 3.1 of [3] asserts that for some $\alpha \in \text{Tor}_q^{UI}(M, \mathbb{k})$, $UL/I \cdot \alpha$ has polynomial growth at least equal to $\dim(L/I)_{\text{even}} - p$. This means that for some positive C ,

$$\sum_{i \leq k} \dim (UL/I \cdot \alpha)_i \geq Ck^{(\dim(L/I)_{\text{even}}) - p}, \quad k \text{ sufficiently large.}$$

Since $\text{Tor}_*^{UI}(M, \mathbb{k})$ is the homology of $\wedge^* sI \otimes M$, it follows that

$$\sum_{i \leq k} \dim \text{Tor}_q^{UI}(M, \mathbb{k})_i \leq \left(\sum_{i \leq k} \dim L_i \right)^q \sum_{i \leq k} \dim M_i.$$

But $(L/I)_{\text{even}} \cong E_{\leq n}$ and so a quick calculation gives

$$\sum_{i \leq k} \dim L_i \geq Kk^{r+1}, \quad k \text{ sufficiently large.} \quad (7)$$

Finally, recall that either $L = L_{\text{even}}$ or else L is generated by the elements of odd degree x_i . In the former case $L = E$ and the Proposition follows from (6) and (7). In the second case we have $L_{\text{odd}} = [x_1, E] + \dots + [x_s, E] + \mathbb{k}x_1 + \dots + \mathbb{k}x_s$, and hence (7) yields

$$\sum_{i \leq k} E_i \geq \frac{K}{s+1} k^{r+1} + s, \quad k \text{ sufficiently large.}$$

Combined with (6) this formula gives the Proposition. $\square$

Proof of Theorem 5: Since L is not solvable we may choose n so that $\dim(L_{\text{even}})_{\leq n} > \text{polydepth } L$ (Theorem 3), and so that the sub Lie algebra generated by $L_{\leq n}$ satisfies $\text{polydepth } E \leq \text{polydepth } L$ (Proposition 1). Then E is not solvable (Proposition 3).

Let $x_1, \dots, x_s$ generate E , and put $d = \max \deg x_i$. Letting UE act via the adjoint representation on E we have that

$$UE_{[0,q]} \cdot E_{[k+1,k+d]} \supset E_{[k+1,k+q]}.$$

For any $r \geq 1$ choose $q = q(r)$ so that $\sum_{i=k+1}^{k+q} \dim E_i \geq k^{r+1}$, k sufficiently large (Proposition 5). Then

$$\sum_{i=k+1}^{k+d} \dim L_i \geq \sum_{i=k+1}^{k+d} \dim E_i \geq \frac{1}{\dim UE_{[0,q]}} k^{r+1} \geq k^r, \quad k \text{ sufficiently large.}$$

Since d is independent of r , the Theorem is proved. $\square$

References

- [1] Y. Felix, S. Halperin, J.-M. Lemaire and J.-C. Thomas, Mod p loop space homology. *Inventiones Math.* **95** (1989), 247-262.
- [2] Y. Felix, S. Halperin, C. Jacobson, C. Löfwall and J.-C. Thomas, The radical of the homotopy Lie algebra. *Amer. J. Math.* **110** (1988), 301-322.
- [3] Y. Felix, S. Halperin and J.-C. Thomas, Lie algebras of polynomial growth. *J. London Math. Soc.* **43** (1991), 556-566.
- [4] Y. Felix, S. Halperin and J.-C. Thomas, Hopf algebras of polynomial growth. *J. Algebra* **125** (1989), 408-417.
- [5] Y. Felix, S. Halperin and J.-C. Thomas, Engel Elements in the Homotopy Lie Algebra. *Journal of Algebra*, **144** (1991), 67-78.
- [6] Y. Felix, S. Halperin and J.-C. Thomas, The category of a map and the grade of a module. *Israel Journal of Mathematics* **78** (1992), 177-196.
- [7] Y. Felix, S. Halperin and J.-C. Thomas, *Growth and Lie brackets in the homotopy Lie algebra*. To appear.
- [8] S. Halperin, Universal enveloping algebra and loop space homology. *J. Pure and Appl. Algebra* **83** (1992), 237-282.
- [9] J.-L. Koszul, Homologie et cohomologie des algèbres de Lie. *Bull. Soc. Math. France* **78** (1950), 65-127.
- [10] J.W. Milnor and J.C. Moore, On the structure of Hopf algebras. *Annals of Math.* **81** (1965), 211-264.

Institut de Mathématiques,
Université Catholique de Louvain
1348 Louvain-La-Neuve, Belgium

College of Computer, Mathematical and Physical Sciences
University of Maryland
College Park, MD 20742-3281, USA

Faculté des Sciences
Université d'Angers
Bd. Lavoisier, 49045 Angers, France